

Incorporating DNA Methyltransferase Inhibitors (DNMTis) in the Treatment of Genitourinary Malignancies – a systematic review

Michal Chovanec^{1,2}, Fadi Taza¹, Maitri Kalra¹, Noah Hahn³, Kenneth P. Nephew⁴, Michael J.

Spinella⁵, Costantine Albany¹

¹ Division of Hematology Oncology, Indiana University Simon Cancer Center, Indianapolis, 535 Barnhill drive, IN, 46202, USA, mchovane@iu.edu, mkalra@iu.edu, ftaza@iu.edu, calbany@iu.edu,

² 2nd Department of Oncology, Faculty of Medicine, Comenius University and National Cancer Institute, Klenova 1, 83310, Bratislava, Slovakia,

³ The Johns Hopkins University School of Medicine, 733 N Broadway, Baltimore, MD, 21205, USA, nhahn4@jhmi.edu,

⁴ Molecular and Cellular Biochemistry Department, Indiana University, Bloomington, 212 S. Hawthorne Drive, IN, 47405, USA, knephew@indiana.edu,

⁵ Department of Comparative Biosciences, the University of Illinois at Urbana-Champaign, 2001 South Lincoln Avenue Urbana, IL 61802, USA, spinella@illinois.edu,

Corresponding Author: Dr. Michal Chovanec, Division of Hematology Oncology, Indiana University Simon Cancer Center, 535 Barnhill drive, 46202, Indianapolis, IN, USA,
^{2nd} Department of Oncology, Faculty of Medicine, Comenius University and National Cancer Institute, Bratislava, Slovakia
Phone: (317) 944-0920, Fax: (317) 944-3646, email: mchovane@iu.edu, twitter: @mikechovanec

Running title: Methyltransferase Inhibitors in Genitourinary Cancer

Funding: This article is supported by Slovak Development and Research Agency under contract

No. APVV-15-0086 for Michal Chovanec

Disclosure: nothing to disclose

Word count: 4 509

This is the author's manuscript of the article published in final edited form as:

Chovanec, M., Taza, F., Kalra, M., Hahn, N., Nephew, K. P., Spinella, M. J., & Albany, C. (2018). Incorporating DNA Methyltransferase Inhibitors (DNMTis) in the Treatment of Genitourinary Malignancies: A Systematic Review. *Targeted Oncology*, 13(1), 49–60. <https://doi.org/10.1007/s11523-017-0546-x>

Abstract

Inhibition of DNA methyltransferases emerged as a novel treatment strategy in solid tumors. Aberrant hypermethylation in promoters of critical tumor suppressor genes created a premise, that treatment with hypomethylating agents may lead to restoration of “normal” epigenome and produce clinically meaningful therapeutic outcome. The aim of this review article is to summarize the current state of knowledge of DNA methyltransferase inhibitors in the treatment of genitourinary malignancies. We performed a literature search of PubMed/Medline, ASCO, ASCO GU and AACR libraries on September 17th 2017. Included reports evaluated pre-clinical and clinical studies assessing the role of DNA methyltransferase inhibitors in genitourinary malignancies. The efficacy of DNA methyltransferase inhibitors in genitourinary malignancies was reported in number of studies and suggested the role of DNA hypomethylation in overcoming resistance to conventional cytotoxic treatments. The clinical significance should be further investigated.

Key points

Increasing knowledge about epigenetic landscape in cancer leads to discovery of promising novel drugs.

Evidence from pre-clinical and clinical studies suggests that DNA methyltransferase inhibitors provide anticancer activity in number of tumors.

DNA methyltransferase inhibitors deliver cytotoxic and chemoresistance overcoming effects in several genitourinary malignancies.

1. Introduction

The conventional paradigm of phenotype manifesting as a sole result of the information encoded in the DNA experienced radical shift over the recent years. Epigenetic changes represent series of mechanisms interfering with the DNA and its function without altering the sequence in coding genes. Both, genetic and epigenetic aberrations cooperate to produce conformational changes in chromatin and biochemically alter gene promoters. Such regulation induces or represses the transcriptional gene activity and may ultimately lead to carcinogenesis ^{1, 2}.

Recent studies indicate that epigenetic silencing may be as important as DNA mutations in tumorigenesis and treatment resistance ³. While mutations represent an irreversible change in the DNA sequence, epigenetic silencing is a reversible process. Known modifications with ability to influence the gene expression without altering the DNA sequence include DNA methylation, histone modifications, nucleosome remodeling induced by ATPases, and regulation via non-coding RNAs³⁻⁶. This review will summarize the current state of pre-clinical and clinical knowledge in targeting DNA methyltransferases, a key messengers for DNA methylation, in genitourinary cancer.

2. Literature search

We performed a literature search of the PubMed/Medline database, and meeting libraries of American Society of Clinical Oncology (ASCO), ASCO Genitourinary Cancers Symposium and

American Association for Cancer Research (AACR) for publications with the terms “epigenetics”, “DNMT”, “DNMTi” “genitourinary”, “testicular cancer”, “germ-cell tumors”, “bladder cancer”, “renal cell carcinoma”, “prostate cancer”, “penile cancer”, “azacitidine”, “decitabine”, “guadecitabine”, “zebularine”, “non-nucleoside”. Combinations of these key words were used for comprehensive search as outlined in Figure 1. The search of literature was performed on September 15th 2017. Original full-text articles published in English were reviewed and the reference lists of key articles were further evaluated. We did not limit our search by the years of publication. Our search was conducted according to the Preferred Reporting Items for Systematic Review and Meta-analysis (PRISMA) statement. Identified reports were reviewed according to the Consolidated Standards of Reporting Trials (CONSORT) criteria. The search resulted in overall 4, 152 publications. One hundred and eleven publications were selected for inclusion in our review article [88 original papers (79%) and 23 (21%) review articles]. The literature search and the inclusion and exclusion criteria are summarized in Figure 1 and Table 1.

3. DNA Methylation

A covalent addition of methyl groups (-CH₃) to the carbon-5 position of cytosine represents a non-coding modification of DNA – a methylation ⁷. DNA methylation at the 5'-CG-3' sequence (also CpG dinucleotide) is a unique, heritable mechanism for regulation of gene transcription. Hypermethylation generally acts as gene silencer by downregulating the transcription of CpG-rich promoter regions (CpG islands) of tumor suppressors ⁸. 5-methylcytosine may undergo a spontaneous deamination to thymine allowing for changes in the chromatin structure. This represents a change in DNA sequence and creates a significant restraint to initiate a transcriptional activity. DNA methylation may be found in distinct locations that may

further influence gene expression and results in different biological phenotypes. Promoter region methylation is a repressive mark, while the gene body methylation is usually associated with active gene expression ^{9, 10}. DNA hypermethylation in the promoter region of tumor suppressor genes in human cancer may represent one of the most important mechanisms in cancer development ¹¹.

DNA methylation was discovered to be present also in context of non-CpG (CpA, CpT, CpC). Non-CpG methylation coexists with CpG methylation, but the exact role is not clear. Evidence suggests an involvement in gene expression when non-CpG methylation is found in promoter regions ¹². Unlike CpG methylation, non-CpG methylation is asymmetrical and thus the same methylation pattern after each cell division cannot be maintained ¹³. As a result, non-CpG methylation is decreased in rapidly proliferating tissues suggesting a potential role in carcinogenesis and pluripotency ^{14, 15}.

The current discoveries have identified significance of enzymatic activity of three DNA methyltransferases (DNMTs): DNMT1, DNMT3A, and DNMT3B. The role of DNMT1 is to maintain the existing methylation patterns, while DNMT3A and DNMT3B are responsible for *de novo* methylation¹⁶.

4. Proposed mechanism of action of DNA methyltransferase inhibitors (DNMTis) in solid tumors

In-depth understanding of the mechanisms underlying the effects of DNMTis is virtually non-existent. Thus, hypotheses and emerging evidence are outlined in this review article in attempt to summarize the current state of knowledge. A well-proven fact that DNMTis are responsible for global hypomethylation resulting in (re)expression of certain tumor suppressor genes offers a superficial knowledge on how DNMTis deliver their effect ^{17, 18}. On the other hand, global

hypomethylation in tumors was linked with aberrant activation of cancer germ-line oncogenes that promote cell proliferation, angiogenesis and metastasis ¹⁹. It is unclear, whether the intra-tumoral hypomethylation activating oncogenes can be overcome by treatment effects of DNMTis which alter the whole body methylome. Non-specificity of treatment with DNMTis may raise doubts about the safety of drugs with the potential to induce re-expression of tumor-suppressor genes as well as oncogenes. Available data from a phase II clinical trial in ovarian cancer did not show any secondary malignancies in patients treated with decitabine for 2-30 months ²⁰. Primary target population for clinical trials evaluating the efficacy of these novel agents are patients with refractory solid tumors pretreated with conventional treatments. Described evidence suggests that survival benefit of treatment with DNMTis in this patient population may outweigh the risk of potential secondary malignancies. The most common toxicities are nausea, constipation, allergic reactions and bone marrow suppression. Grade 3-4 neutropenia and thrombocytopenia were seen in 22% and 11% of patients respectively ²⁰.

Most of the initial clinical trials using a single agent DNMTi to treat solid tumors were unsuccessful in improving outcomes. DNMTis were used at high doses with the intent to deliver a non-specific cytotoxic effect, however, the delivery at low doses are optimized to inhibit DNA methylation ²⁰⁻²². More recently, trials are testing whether DNMTis could sensitize refractory tumors to chemotherapy and whether they can improve immune response and boost immune therapy.

4.1 DNMTis to overcome chemo-resistance

Multiple hypotheses are proposed to explain the role of DNMTis in overcoming resistance to chemotherapy, specifically the DNA targeted drugs (DTDs). Lethal DNA damage is the main mechanism of cell death induced by platinum compounds and doxorubicin ²³. Cisplatin induces

DNA breaks by creating inter-strand adducts. Doxorubicin also creates DNA damage after being incorporated into DNA. DNMTis may potentiate the DNA damage induced by cisplatin and doxorubicin by increasing the accessibility of these drugs through loosening chromatin globally, which is required for DTD incorporation and DNA damage²⁴. Furthermore, epigenetic synergy of decitabine and platinum agents, but no other cytotoxic drugs, was observed by Qin *et al.* in colon cancer cell line. Treating the cell line with 16 cytotoxic drugs including platinum agents did not result in activation of hypermethylated cytomegalovirus promoter. However, the addition of platinum compounds achieved a striking synergy in activating the promoter. Experiments also resulted into significantly better reactivation of hypermethylated tumor suppressor genes (*MHLMI* and *PDLIM4*) with the combination of decitabine and carboplatin, compared to each drug alone, therefore offering evidence that platinum resistance can be overcome with the addition of decitabine¹⁸

4.2 DNA methyltransferase (DNMT) inhibitors and immune responses

DNMTis have potency to reactivate the silenced tumor suppressor genes by DNMT inhibition, and are capable of upregulating the genes encoding MHC I molecules, tumor antigens²⁵⁻²⁷ and interferon (IFN) response proteins²⁸. Decitabine upregulated chemokine expression *in vitro* in ovarian tumor cell and increased the number of NK and CD8+ cells in malignant ascites in an orthotopic mouse model²⁹.

Recent reports suggest immune modulation as a possible mechanism of action for azacitidine. The expression of endogenous retroviruses (ERVs) is silenced by DNA methylation^{30,31}, and treatment with low doses of azacitidine reactivated ERVs in ovarian and colon cancer cells^{32,33}. While most ERVs have lost the ability to mature and infect other cells, they can still expand within their host

genome^{27, 34}. Upon the expression of ERV, a double-strand RNA (dsRNA) that is constructed plays a role in creating of a type I IFN response^{32, 33}. As a result, azacitidine may induce tumor cells to mimic virally infected cells, resulting in an antitumor immune response³⁵. Several clinical trials with azacitidine or guadecitabine in combination with immune checkpoint inhibitors are ongoing in various solid tumors.

5. DNA methyltransferase inhibitors (DNMTis) and their mechanism of action

Historical pre-clinical and clinical studies assessed 5-azacitidine (azacitidine) and 5-aza-2'-deoxycytidine (decitabine) that were among the first agents efficient in downregulating the DNA methylation. Azacitidine and decitabine act as inhibitors of DNMTs and are now referred to as “first generation DNMTis” preceding a “second generation DNMTis” guadecitabine (2'-deoxy-5-azacytidyl-(3'-5')-2'-deoxyguanosine sodium salt) and zebularine (1-[beta]-D-ribofuranosyl]-1,2-dihydropyrimidin-2-one). These agents act as cytidine antimetabolite analogues creating covalent complexes with DNMTs and DNA. Trapping the DNMTs in this fashion ultimately alters their function resulting in downregulation of DNA methylation³⁶. This covalent trapping, however, also induces DNA damage that was suggested in the mutagenic and cytotoxic effects of DNMTis³⁷. Azacitidine and decitabine were therefore initially developed as chemotherapeutic agents and were used at cytotoxic doses.

First generation DNMTis suffer from fast metabolism induced by hydrolytic cleavage and deamination. As a result, their stability in the blood is limited and their use as cytotoxic agents in solid cancers proved unsuccessful³⁸. The second-generation DNMTis guadecitabine and zebularine are stable in the liquid environment and allow for a longer effective half-life³⁸⁻⁴⁰. While hypomethylation induced by the first generation DNMTis in *in-vitro* studies⁴¹⁻⁴³ brings evidence

of effectivity of these agents, downregulating the DNA methylation with guadecitabine in cancer xenograft models ⁴⁴ provided proof that unfavorable pharmacokinetic profile of first generation DNMTis can be overcome. Zebularine was shown to be preferentially incorporated in zebularine-sensitive tumor cells with specific transcriptomic and epigenomic signature in *in-vitro* and xenograft mouse model experiments with human liver cancer. This shows a promising tool for predicting response to this second generation DNMTi ⁴⁵. Fluorocyclo-pentenylcytosine is another novel next generation nucleoside agent with antimetabolic activity that also acts as DNMT1 inhibitor. Numerous preclinical cell-line and xenograft animal models have shown antitumor activity of this drug ⁴⁶⁻⁴⁹.

Alternative mechanism that reduces CpG island methylation is inhibition of DNMT1 with antisense oligonucleotides which are not incorporated into genomic DNA. In-vitro experiments in bladder cancer cells have shown that MG88 targeting the 3'-untranslated region of *DNMT1* resulted in suppressed DNMT1 expression, thus allowing for re-expression of tumor suppressor gene *α-CDKN2A* ¹⁷.

Mechanism involved in transport of DNMTis across cell membrane indirectly influence the ability of DNMTis to perform effectively. Human organic cation and nucleoside transporters may both mediate intake or efflux of azacitidine, decitabine and zebularine. These transporters, thus, contribute to chemoresistance or chemosensitivity to DNMTis in cancer therapy ⁵⁰.

Several other non-nucleoside targeting DNMTi agents are procaine ⁵¹ N-acetylprocainamide, procainamide (perturbing the interactions between the protein and its target sites), hydralazine (decreases the expression of DNMT1 and 3A)⁵², (–)-epigallocatechin-3-gallate (EGCG, a catalytic pocket blocker of DNMT1 found in green tea)⁵³, and RG108 (the first rationally designed inhibitor of DNMTs)⁵⁴. Non-nucleoside agents have shown considerably less demethylating activity in

bladder and prostate cancer cell lines compared to decitabine⁵². Ongoing clinical trials using DNMTi agents are summarized in Table 2.

6. Targeting DNMTs in genitourinary cancer

Following chapters are discussing a current state of evidence in the pre-clinical and clinical use of DNMTis in genitourinary malignancies.

6.1 Germ Cell Tumors

Germ cell tumors (GCTs) have a global DNA hypomethylation that may explain an exceptional sensitivity to platinum-based chemotherapy⁵⁵⁻⁵⁷ resulting from the highest de novo DNA methyltransferase expression (DNMT3A/3B) among solid cancers (Figure 2)^{58, 59}. However, numerous promoters in GCT cell lines were discovered to be hypermethylated in non-seminomas⁶⁰ and promoter hypermethylation of *RASSF1A* and *HIC1* genes was linked to cisplatin resistance in embryonal carcinoma cell lines⁶¹. Beyrouthy *et al.* have shown that overexpression of DNMT3B is associated with hypersensitivity to decitabine. Treatment with decitabine resulted in a re-sensitization of testicular cancer cells to cisplatin. Furthermore, the demethylation resulted in a re-activation of tumor suppressor genes⁶². Similar observations were made by Wermann *et al.*, who observed an increased sensitivity to cisplatin after treatment with 5-azacitidine in platinum resistant GCT cell lines⁵⁷.

We evaluated guadecitabine *in vitro* and in an *in vivo* in a mouse model of cisplatin refractory GCT and found that testicular cancer cells had exquisite sensitivity to single agent exposure. Guadecitabine completely abolished progression and induced complete regression of cisplatin resistant testicular cancer xenografts even at doses well below those required to impact somatic

solid tumors ²². A phase I study of guadecitabine in combination with cisplatin in refractory GCT subjects who relapsed after multiple lines of chemotherapy including high-dose chemotherapy (HDCT) is currently ongoing at Indiana University (NCT02429466). Anecdotal evidence from a phase II study evaluating hydralazine and magnesium valproate showed a stable disease in 1 patient with refractory non-seminoma ⁶³.

6.2 Bladder cancer

Aberrant DNA methylation was initially found to be relevant to carcinogenesis in human bladder tumors and cell lines. Later, comprehensive genomic and promoter assessments have shown characteristic DNA methylation patterns in bladder cancer ^{64, 65}. Maruyama *et al.* assessed the promoter methylation status of several cancer-related genes including *CDH1*, *RASSF1A*, *APC*, *CDH13*, *FHIT*, *RARβ*, *GSTP1*, *p16INK4a*, *DAPK* and *MGMT* in 98 bladder tumors. High methylation frequency in *RASSF1A*, *APC*, *CDH1*, *CDH13* and *FHIT* correlated with poor prognostic clinical-pathological features and shorter overall survival ⁶⁶. Hypermethylation of *RASSF1A*, *APC* and *MGMT* has been linked to high grade and invasive tumors in 76 patients with urothelial bladder cancer ⁶⁷. In pre-clinical studies, azacitidine inhibited the proliferation of bladder cancer cells and arrested cells at the G0/G1 phase. Wang *et al.* performed *in vivo* and *in vitro* experiments showing that azacitidine markedly downregulated the expression of DNMT3A/3B, reactivated expression of hepaCAM, and inhibited bladder cancer growth in nude mice ⁶⁸. Treatment with demethylating agents decitabine and zebularine (nucleoside analog of cytidine) induced a growth inhibition in urothelial bladder cancer and renal cell cancer cells resulting in a 17-132% prolongation of cell doubling time ⁶⁹. A preclinical study evaluating single agent azacitidine in 19 dogs with spontaneous urothelial bladder cancer confirmed a

myelosuppressive, but relatively safe toxicity profile. Partial remission was seen in 22%, stable disease in 50% and progressive disease in 22% of dogs evaluable for tumor response ⁷⁰. Another pre-clinical study showed that sensitization of cisplatin resistant bladder cancer cell lines can be achieved *in vitro* by decitabine pre-treatment alone and in combination with the HDAC inhibitor vorinostat ⁷¹. Chuang *et al.* conducted a pre-clinical *in vivo* assessment of single agent decitabine and guadecitabine in murine xenograft models derived from bladder cancer cell lines. Intraperitoneal delivery of both drugs was effective in reducing the level of DNA methylation at the P16 promoter. Murine tumors also exhibited a growth inhibition, although this treatment was insufficient to reduce the size of the tumors. A subcutaneous administration delivered similar results. Toxicity profile, however, was more favorable with subcutaneous injection ⁷². While zebularine effectively induced hypomethylation⁷³ and reactivated silenced genes⁷⁴ in T24 bladder cell lines, another study showed its' complex metabolism and limited incorporation into the DNA in bladder cancer cell lines. This may be the reason for lower efficacy of zebularine compared to azacitidine and decitabine in bladder cancer *in vitro* ⁷⁵. Interestingly, zebularine increased radiation-induced DNA damage in bladder cancer cell line and increased the radiation induced tumor growth delay in xenograft mouse model ⁷⁶. The function of DNMT1 in bladder cancer cell lines was recently discovered to be mediated by the virtue of long non-coding (lnc) RNA *DBCCR1-003* derived from the locus of *DBCCR1* tumor suppressor gene ⁷⁷. Habuchi *et al.* previously linked the loss of *DBCCR1* expression and its association with transitional-cell bladder cancer ⁷⁸. Qi *et al.* reported that *DBCCR1-003* normally binds to DNMT1 and prevents the hypermethylation of *DBCCR1* thus allowing its expression. Treatment with decitabine or overexpression of *DBCCR1-003* resulted into increased expression of *DBCCR1* via reversed promoter hypermethylation and DNMT1 binding to *DBCCR1-003* and promoter *DBCCR1* in T24

bladder cancer cell line. Such process led to significant growth inhibition of the cell line suggesting DBCCR1-003 as novel biomarker and potential treatment target ⁷⁷. *In-vivo* experiments in patient-derived bladder cancer xenografts in mice showed that guadecitabine reduced DNA methylation at the *p16* promoter region and reduced the growth of tumors in xenografts ⁷⁹.

A combination of azacitidine with sodium phenylbutyrate was used in a phase I study of patients with refractory solid tumors. Two of 28 patients had bladder cancer, however, no objective responses were observed within this trial ⁸⁰. Another phase I study used azacitidine with valproic acid in 55 patients with refractory malignancies of whom 3 had advanced urothelial cancer. The combination did not produce any objective responses. Stable disease was achieved in 25% of patients with various malignancies, other than urothelial carcinoma ⁸¹.

Two clinical trials evaluating treatment with decitabine (NCT00030615) and CC-486, oral formulation of azacitidine (NCT01478685) in solid tumors including bladder cancer were conducted, but to our best knowledge, the results were not reported yet. Another clinical study (NCT02223052) with CC-486 in solid tumors including genitourinary cancer is currently underway. Additionally, a genomic-based assignment of treatment including azacitidine and decitabine is currently ongoing in advanced urothelial carcinoma (NCT02788201).

A phase I study of antisense oligonucleotide inhibitor of DNMT1 assessed the safety and efficacy of MG98, given in an infusion over 7 days to 33 patients with solid tumors, including bladder, upper urinary tract and prostate cancer. Treatment related toxicities were generally mild and most common being fatigue, headache and myalgia of grade ≤ 2 . Dose limiting toxicities were grade 3 transaminitis and grade 3 thrombocytopenia. Evidence of activity was observed in this study, however, none of the patients that responded had genitourinary cancers ⁸².

Fluorocyclo-pentenylcytosine (RX-3117) have shown antitumor activity in gemcitabine resistant pre-clinical models and is currently being evaluated in a phase I/II study in pancreatic and urothelial cancer (NCT02030067) ⁴⁶⁻⁴⁹.

The insufficient clinical activity of most of the DNMTi in urothelial cancer could be explained by the short half-life of the first generation DNMTi agents. From a mechanistic standpoint, DNMTi may be better used in combination with cytotoxic drugs such as cisplatin or with immune checkpoint inhibitors.

6.3. Renal Cell Carcinoma

Abnormal hypermethylation as well as hypomethylation of DNA may occur in renal cell carcinoma (RCC), resulting in activation of tumorigenesis and chromosomal instability ^{83, 84}. Numerous tumor suppressor genes have been reported to be partially or completely silenced due to hypermethylation of their enhancer and promoters leading to increased tumor cell proliferation, invasion, and metastasis ⁸⁵. Deep DNA methylation and transcriptome profiling of diverse histological RCC subtypes uncovered that clear cell (ccRCC), papillary and translocation RCC as well as mucinous and spindle cell carcinomas, are 3-fold more hypermethylated as compared to oncocytoma and chromophobe RCC ⁸⁶. Moris *et al.* described nine genes showing frequent promoter region methylation in primary RCC tumor samples. The methylation of SCUBE3 was associated with a significantly increased risk of cancer death or relapse ⁸⁷. Li *et al.* found that DNMT1 protein was expressed significantly higher in ccRCC compared to normal tissues (56.2% and 27.3%, respectively). The expression of DNMT1 was positively correlated with tumor size, highly malignant phenotype, lymph node metastasis, vascular invasion, recurrence and poor prognosis. This observation was confirmed *in vitro* in cell lines, where knockdown of DNMT1 significantly inhibited ccRCC cell viability, induced apoptosis, decreased colony formation and

the invading ability ⁸⁸. German group recently proposed metastasis associated methylome signature obtained from genome-wide TCGA dataset. Authors predicted metastatic disease with 93% sensitivity and 89% specificity and proposed the prospective validation of this tool ⁸⁹.

Treatment with azacitidine resulted in suppression of cell proliferation in all 15 RCC cell lines evaluated by Ricketts *et al.* Interestingly, response correlated with alterations in VHL promoter methylation, however some cell lines without VHL tumor suppressor gene methylation responded to treatment as well. This finding is suggestive of other hypermethylated suppressor genes activated with DNMTi treatment with several hypermethylated candidate tumor suppressor genes identified (RGS7, NEFM, TMEM74, GCM2 and AEBP1). Methylation of GCM2, NEFM and RGS7 also strongly correlated with poor prognosis ⁹⁰. Treatment of A-498 RCC cell line with low-dose zebularine resulted in limited cell inhibition, however, authors observed an up-regulation and down-regulation of 308 and 253 gene transcripts, respectively. Many of the re-expressed genes belong to metallothionein family, potent protectors against oxidative stress, that were discovered to be down-regulated in RCC tumors ⁹¹. Decitabine, but not zebularine was able to reexpress a hypermethylated VHL gene in RCC cell lines and caused a tumor shrinkage in RCC xenograft mouse model. Only tumors with hypermethylated VHL gene responded to treatment. VHL mutated mice did not show any response ⁹². Guadecitabine showed an interesting potential to increase immunogenicity of renal cell carcinoma and other tumors cell lines. Treatment with guadecitabine induced a de novo expression or reexpression of cancer testis antigen related genes (*MAGE-A1*, *MAGE-A2*, *MAGE-A3*, *MAGE-A4*, *MAGE-A10*, *GAGE 1-2*, *GAGE 1-6*, *NY-ESO-1*, and *SSX 1-5*) and HLA class I antigens of ICAM-1, an important factor for improved recognition of cancer cells by gp-100 specific cytotoxic T lymphocytes ⁹³.

Reu and colleagues have shown a synergistic potential of cytokine therapy combined with decitabine. Resistance to the antiproliferative and apoptotic effects of interferons (IFNs) was postulated to result from silencing of IFN response genes by promoter hypermethylation. Treatment of human RCC cells with decitabine overcame IFN resistance, while normal epithelial kidney cells remained unaffected. IFN response gene expression was augmented greater than 10 times by decitabine ⁹⁴.

A phase I study conducted by Lin *et al.* aimed to find the safe and effective dose of azacitidine in combination sodium phenylbutyrate in refractory solid tumors. Three of patients on this study had renal cell carcinoma. While clinical responses were disappointing, treatment induced undetectable activity of DNMTs in renal cell carcinoma patients at low doses (15mg/m²) ⁸⁰. An ongoing phase I/II clinical trial is assessing the feasibility, safety, and efficacy of anti-PD-1 antibody in combination with low-dose decitabine in patients with relapsed or refractory malignancies including RCC (NCT02961101). Another phase I/II trial of azacitidine in combination with bevacizumab for advanced RCC previously treated with VEGF- or mTOR-targeted therapy with or without prior immunotherapy was recently concluded and the results are pending (NCT00934440).

6.4 Prostate Cancer

Among all solid tumors evaluated in TCGA, prostate cancer has the lowest level of expression of *DNMT3A/B* (Figure 2). *GSTP1* is, however, frequently methylated gene in prostate cancer ⁹⁵. Functional epigenetic analyses in prostate cancer cell lines showed the reexpression of genes regulated by hypermethylation their promoters after treatment with zebularine or decitabine (*IFI6*, *GSTP1*) ⁹⁵⁻⁹⁷. While zebularine failed to achieve the reexpression of *GCTP1* it effectively

reexpressed two GST detoxifying enzymes (GST-pi and GST-mu) in mice ^{96, 97}. Another commonly hypermethylated promoter in *ASC* gene (apoptosis associated speck-like protein) was demethylated and reexpressed by decitabine and zebularine in five prostate cancer cell lines ⁹⁸. Gertych *et al.* demonstrated that treatment with azacitidine and zebularine changed the DNA topology status in terms of DNA-histone complex decondensation, along with demethylating effects in prostate and liver cancer cells ⁹⁹. A study by Kim *et al.* described that overexpression of lysine-specific demethylase 4A (JMJD2A) resulted in initiation of prostate cancer development in mice further mediated by JMJD2A/ETV1/YAP1 pathway ¹⁰⁰. In further preclinical studies, azacitidine, demonstrated synergistic effects with docetaxel and cisplatin in androgen receptor (AR)-positive 22RV1 and in AR-negative PC3 cells ¹⁰¹. Decitabine also exhibited synergy with cisplatin and cyclophosphamide in non-prostate cancer cell lines, although the relationship to induced DNA demethylation was unclear ¹⁰². Prostate cancer cell lines that were treated with decitabine showed partial demethylation of *TMSI/ASC* locus (a frequently hypermethylated gene in prostate cancer) and the subset of alleles remained unmethylated for over 3 months while others were remethylated within 1 week ¹⁰³. Azacitidine prevented *de novo* development of cancer in all of 14 mice used in experiments with the model of transgenic adenocarcinoma of the mouse prostate ¹⁰⁴.

Azacitidine and sodium phenylbutyrate failed to induce the response in a phase I trial including 5 prostate cancer patients, perhaps owing to the undetectable DNMT activity before the treatment ⁸⁰. Another phase I trial evaluating azacitidine and valproic acid where 2 patients with advanced prostate cancer were included reported stable disease in 1 patient ⁸¹. In a phase II trial, subcutaneous azacitidine did not resensitize tumors to androgen-deprivation therapy in 36 patients with progressive castration resistant prostate cancer (CRPC) ¹⁰⁵. However, the combination of

azacitidine with docetaxel was active in metastatic CRPC patients ¹⁰⁶. In a phase I/II study, azacitidine was given daily for 5 days followed by docetaxel on day 6. PSA response was seen in 10 of 19 evaluable patients and objective response was observed in 3 of 10 evaluable patients. Significant demethylation of GADD45A was observed with azacitidine treatment ¹⁰⁶.

Thus far, the clinical efficacy outcomes of azacitidine in prostate cancer trials have demonstrated only modest activity. Possible reasons may be the low level of DNMT3 expression or instability of DNA methylation inhibitors due to the short half life ¹⁰⁷. Wong *et al.* provided strong evidence for DNA methylation recovery and found that H3K9 trimethylation and H3K27 trimethylation were closely associated with DNA methylation recovery ¹⁰⁸. Overall, the efficacy of DNMTi in prostate cancer treatment is yet to be determined ².

6.5 Penile cancer

Several studies summarized by Kuasne et al. attempted to describe epigenetic alterations in penile carcinoma (PeCa). Hypermethylation of the CDK2A gene promoter was found to be present in 15 to 42% of PeCa samples ¹⁰⁹. Feber et al. evaluated the methylation profile of 38 PeCa samples using high-density genome-wide methylation arrays. The authors identified a clear hypermethylation profile associated with cancer phenotype and identified novel epigenetic signatures associated with HPV infection and loco-regional spread. Interestingly, epigenetic signatures that were predictive of metastases in lymph nodes suggested a lower metastatic potential of hypermethylated tumors. In addition, HPV infection showed significant correlation with DNA methylation. The majority of HPV-positive samples were hypomethylated and showed better clinical outcome ¹¹⁰. Distinct methylome and transcriptome patterns were described in more recent genome-wide methylation and transcriptome analysis. Aberrant DNA methylation was linked to

expression of specific genes connected to higher tumor aggressiveness and shorter duration of survival ¹¹¹. These studies suggest a role for DNA methylation in PeCa. Nevertheless, to our knowledge pre-clinical and clinical trials with DNMTis have not been reported.

Conclusion

Epigenetic targeting is an exciting new field in cancer research. Pre-clinical efforts to elucidate underlying mechanisms of treatment resistance have led to the initiation of several clinical trials involving epigenetic modulation in genitourinary malignancies. Current knowledge does not yet robustly support the immediate incorporation of DNMTis into treatment of genitourinary cancer. However, DNA methylation appears to be an important mechanism of treatment resistance, which may be overcome by incorporating DNMTis into the therapy. Initial studies evaluating older generations of DNMTis have shown insufficient activity due to unfavorable pharmacokinetics. Nevertheless, the discovery of new generation of DNMTis, coupled to a better understanding of mechanisms of action, have provided the rationale for combination therapies, which may lead to more favorable clinical outcomes. Large clinical studies are needed to provide better understanding whether these agents will find place in everyday treatment of genitourinary cancer.

Compliance with ethical standards

This article is supported by Slovak Development and Research Agency under contract No. APVV-15-0086 for Michal Chovanec. All authors declare no conflict of interest.

References

1. Hanahan D, Weinberg RA. Hallmarks of cancer: the next generation. *Cell* 2011; 144:646-74.
2. Albany C, Alva AS, Aparicio AM, Singal R, Yellapragada S, Sonpavde G, Hahn NM. Epigenetics in prostate cancer. *Prostate Cancer* 2011; 2011:580318.

3. Herman JG, Baylin SB. Gene silencing in cancer in association with promoter hypermethylation. *The New England journal of medicine* 2003; 349:2042-54.
4. Becker PB, Workman JL. Nucleosome remodeling and epigenetics. *Cold Spring Harb Perspect Biol* 2013; 5.
5. Grunstein M. Histone acetylation in chromatin structure and transcription. *Nature* 1997; 389:349-52.
6. Eddy SR. Non-coding RNA genes and the modern RNA world. *Nat Rev Genet* 2001; 2:919-29.
7. Jones PA, Baylin SB. The epigenomics of cancer. *Cell* 2007; 128:683-92.
8. Das PM, Singal R. DNA methylation and cancer. *Journal of clinical oncology : official journal of the American Society of Clinical Oncology* 2004; 22:4632-42.
9. Takai D, Jones PA. Comprehensive analysis of CpG islands in human chromosomes 21 and 22. *Proceedings of the National Academy of Sciences of the United States of America* 2002; 99:3740-5.
10. Jeschke J, Collignon E, Fuks F. DNA methylome profiling beyond promoters - taking an epigenetic snapshot of the breast tumor microenvironment. *FEBS J* 2015; 282:1801-14.
11. Yan W, Herman JG, Guo M. Epigenome-based personalized medicine in human cancer. *Epigenomics* 2016; 8:119-33.
12. Lister R, Pelizzola M, Dowen RH, Hawkins RD, Hon G, Tonti-Filippini J, Nery JR, Lee L, Ye Z, Ngo QM, et al. Human DNA methylomes at base resolution show widespread epigenomic differences. *Nature* 2009; 462:315-22.
13. Patil V, Ward RL, Hesson LB. The evidence for functional non-CpG methylation in mammalian cells. *Epigenetics* 2014; 9:823-8.
14. Han H, Cortez CC, Yang X, Nichols PW, Jones PA, Liang G. DNA methylation directly silences genes with non-CpG island promoters and establishes a nucleosome occupied promoter. *Hum Mol Genet* 2011; 20:4299-310.
15. Ziller MJ, Muller F, Liao J, Zhang Y, Gu H, Bock C, Boyle P, Epstein CB, Bernstein BE, Lengauer T, et al. Genomic distribution and inter-sample variation of non-CpG methylation across human cell types. *PLoS Genet* 2011; 7:e1002389.
16. Kim H, Park J, Jung Y, Song SH, Han SW, Oh DY, Im SA, Bang YJ, Kim TY. DNA methyltransferase 3-like affects promoter methylation of thymine DNA glycosylase independently of DNMT1 and DNMT3B in cancer cells. *Int J Oncol* 2010; 36:1563-72.
17. Fournel M, Sapieha P, Beaulieu N, Besterman JM, MacLeod AR. Down-regulation of human DNA-(cytosine-5) methyltransferase induces cell cycle regulators p16(ink4A) and p21(WAF/Cip1) by distinct mechanisms. *The Journal of biological chemistry* 1999; 274:24250-6.
18. Qin T, Si J, Raynal NJ, Wang X, Gharibyan V, Ahmed S, Hu X, Jin C, Lu Y, Shu J, et al. Epigenetic synergy between decitabine and platinum derivatives. *Clin Epigenetics* 2015; 7:97.
19. Van Tongelen A, Lorient A, De Smet C. Oncogenic roles of DNA hypomethylation through the activation of cancer-germline genes. *Cancer Lett* 2017; 396:130-7.
20. Matei D, Fang F, Shen C, Schilder J, Arnold A, Zeng Y, Berry WA, Huang T, Nephew KP. Epigenetic resensitization to platinum in ovarian cancer. *Cancer research* 2012; 72:2197-205.
21. Tsai HC, Li H, Van Neste L, Cai Y, Robert C, Rassool FV, Shin JJ, Harbom KM, Beaty R, Pappou E, et al. Transient low doses of DNA-demethylating agents exert durable antitumor effects on hematological and epithelial tumor cells. *Cancer Cell* 2012; 21:430-46.
22. Albany C, Hever-Jardine MP, von Herrmann KM, Yim CY, Tam J, Warzecha JM, Shin L, Bock SE, Curran BS, Chaudhry AS, et al. Refractory testicular germ cell tumors are highly sensitive to the second generation DNA methylation inhibitor guadecitabine. *Oncotarget* 2017; 8:2949-59.
23. Bouwman P, Jonkers J. The effects of deregulated DNA damage signalling on cancer chemotherapy response and resistance. *Nat Rev Cancer* 2012; 12:587-98.

24. Fang F, Munck J, Tang J, Taverna P, Wang Y, Miller DF, Pilrose J, Choy G, Azab M, Pawelczak KS, et al. The novel, small-molecule DNA methylation inhibitor SGI-110 as an ovarian cancer chemosensitizer. *Clinical cancer research : an official journal of the American Association for Cancer Research* 2014; 20:6504-16.
25. Karpf AR, Lasek AW, Ririe TO, Hanks AN, Grossman D, Jones DA. Limited gene activation in tumor and normal epithelial cells treated with the DNA methyltransferase inhibitor 5-aza-2'-deoxycytidine. *Mol Pharmacol* 2004; 65:18-27.
26. Karpf AR, Peterson PW, Rawlins JT, Dalley BK, Yang Q, Albertsen H, Jones DA. Inhibition of DNA methyltransferase stimulates the expression of signal transducer and activator of transcription 1, 2, and 3 genes in colon tumor cells. *Proc Natl Acad Sci U S A* 1999; 96:14007-12.
27. Gotwals P, Cameron S, Cipolletta D, Cremasco V, Crystal A, Hewes B, Mueller B, Quarantino S, Sabatos-Peyton C, Petruzzelli L, et al. Prospects for combining targeted and conventional cancer therapy with immunotherapy. *Nat Rev Cancer* 2017; 17:286-301.
28. Li H, Chiappinelli KB, Guzzetta AA, Easwaran H, Yen RW, Vatapalli R, Topper MJ, Luo J, Connolly RM, Azad NS, et al. Immune regulation by low doses of the DNA methyltransferase inhibitor 5-azacitidine in common human epithelial cancers. *Oncotarget* 2014; 5:587-98.
29. Wang L, Amoozgar Z, Huang J, Saleh MH, Xing D, Orsulic S, Goldberg MS. Decitabine Enhances Lymphocyte Migration and Function and Synergizes with CTLA-4 Blockade in a Murine Ovarian Cancer Model. *Cancer Immunol Res* 2015; 3:1030-41.
30. Dewannieux M, Heidmann T. Endogenous retroviruses: acquisition, amplification and taming of genome invaders. *Curr Opin Virol* 2013; 3:646-56.
31. Kassiotis G, Stoye JP. Immune responses to endogenous retroelements: taking the bad with the good. *Nat Rev Immunol* 2016; 16:207-19.
32. Chiappinelli KB, Strissel PL, Desrichard A, Li H, Henke C, Akman B, Hein A, Rote NS, Cope LM, Snyder A, et al. Inhibiting DNA Methylation Causes an Interferon Response in Cancer via dsRNA Including Endogenous Retroviruses. *Cell* 2015; 162:974-86.
33. Roulois D, Loo Yau H, Singhania R, Wang Y, Danesh A, Shen SY, Han H, Liang G, Jones PA, Pugh TJ, et al. DNA-Demethylating Agents Target Colorectal Cancer Cells by Inducing Viral Mimicry by Endogenous Transcripts. *Cell* 2015; 162:961-73.
34. Thompson PJ, Macfarlan TS, Lorincz MC. Long Terminal Repeats: From Parasitic Elements to Building Blocks of the Transcriptional Regulatory Repertoire. *Mol Cell* 2016; 62:766-76.
35. Terranova-Barberio M, Thomas S, Munster PN. Epigenetic modifiers in immunotherapy: a focus on checkpoint inhibitors. *Immunotherapy* 2016; 8:705-19.
36. Weisenberger DJ, Velicescu M, Cheng JC, Gonzales FA, Liang G, Jones PA. Role of the DNA methyltransferase variant DNMT3b3 in DNA methylation. *Mol Cancer Res* 2004; 2:62-72.
37. Juttermann R, Li E, Jaenisch R. Toxicity of 5-aza-2'-deoxycytidine to mammalian cells is mediated primarily by covalent trapping of DNA methyltransferase rather than DNA demethylation. *Proceedings of the National Academy of Sciences of the United States of America* 1994; 91:11797-801.
38. Orta ML, Pastor N, Burgos-Moron E, Dominguez I, Calderon-Montano JM, Huertas Castano C, Lopez-Lazaro M, Helleday T, Mateos S. Zebularine induces replication-dependent double-strand breaks which are preferentially repaired by homologous recombination. *DNA Repair (Amst)* 2017; 57:116-24.
39. Yoo CB, Jeong S, Egger G, Liang G, Phiasivongsa P, Tang C, Redkar S, Jones PA. Delivery of 5-aza-2'-deoxycytidine to cells using oligodeoxynucleotides. *Cancer Res* 2007; 67:6400-8.
40. Yoo CB, Jeong S, Egger G, Liang G, Phiasivongsa P, Tang C, Redkar S, Jones PA. Delivery of 5-Aza-2'-Deoxycytidine to Cells Using Oligodeoxynucleotides. *Cancer research* 2007; 67:6400-8.
41. Cho M, Grabmaier K, Kitahori Y, Hiasa Y, Nakagawa Y, Uemura H, Hirao Y, Ohnishi T, Yoshikawa K, Oosterwijk E. Activation of the MN/CA9 gene is associated with hypomethylation in human renal cell carcinoma cell lines. *Mol Carcinog* 2000; 27:184-9.

42. Nguyen CT, Weisenberger DJ, Velicescu M, Gonzales FA, Lin JC, Liang G, Jones PA. Histone H3-lysine 9 methylation is associated with aberrant gene silencing in cancer cells and is rapidly reversed by 5-aza-2'-deoxycytidine. *Cancer research* 2002; 62:6456-61.
43. Biswal BK, Beyrouthy MJ, Hever-Jardine MP, Armstrong D, Tomlinson CR, Christensen BC, Marsit CJ, Spinella MJ. Acute hypersensitivity of pluripotent testicular cancer-derived embryonal carcinoma to low-dose 5-aza deoxycytidine is associated with global DNA Damage-associated p53 activation, anti-pluripotency and DNA demethylation. *PloS one* 2012; 7:e53003.
44. Sonnenburg D, Spinella MJ, Albany C. Epigenetic Targeting of Platinum Resistant Testicular Cancer. *Curr Cancer Drug Targets* 2015.
45. Andersen JB, Factor VM, Marquardt JU, Raggi C, Lee YH, Seo D, Conner EA, Thorgeirsson SS. An integrated genomic and epigenomic approach predicts therapeutic response to zebularine in human liver cancer. *Science translational medicine* 2010; 2:54ra77.
46. Zhao LX, Yun M, Kim HO, Lee JA, Choi WJ, Lee KM, Lee SK, Lee YB, Ahn CH, Jeong LS. Design, synthesis, and anticancer activity of fluorocyclopentenyl-pyrimidines. *Nucleic Acids Symp Ser (Oxf)* 2005:107-8.
47. Jeong LS, Zhao LX, Choi WJ, Pal S, Park YH, Lee SK, Chun MW, Lee YB, Ahn CH, Moon HR. Synthesis and antitumor activity of fluorocyclopentenyl-pyrimidines. *Nucleosides Nucleotides Nucleic Acids* 2007; 26:713-6.
48. Choi WJ, Chung HJ, Chandra G, Alexander V, Zhao LX, Lee HW, Nayak A, Majik MS, Kim HO, Kim JH, et al. Fluorocyclopentenyl-cytosine with broad spectrum and potent antitumor activity. *Journal of medicinal chemistry* 2012; 55:4521-5.
49. Peters GJ, Smid K, Vecchi L, Kathmann I, Sarkisjan D, Honeywell RJ, Losekoot N, Ohne O, Orbach A, Blaugrund E, et al. Metabolism, mechanism of action and sensitivity profile of fluorocyclopentenylcytosine (RX-3117; TV-1360). *Investigational new drugs* 2013; 31:1444-57.
50. Arimany-Nardi C, Errasti-Murugarren E, Minuesa G, Martinez-Picado J, Gorboulev V, Koepsell H, Pastor-Anglada M. Nucleoside transporters and human organic cation transporter 1 determine the cellular handling of DNA-methyltransferase inhibitors. *British journal of pharmacology* 2014; 171:3868-80.
51. Villar-Garea A, Fraga MF, Espada J, Esteller M. Procaine is a DNA-demethylating agent with growth-inhibitory effects in human cancer cells. *Cancer research* 2003; 63:4984-9.
52. Chuang JC, Yoo CB, Kwan JM, Li TW, Liang G, Yang AS, Jones PA. Comparison of biological effects of non-nucleoside DNA methylation inhibitors versus 5-aza-2'-deoxycytidine. *Mol Cancer Ther* 2005; 4:1515-20.
53. Fang MZ, Wang Y, Ai N, Hou Z, Sun Y, Lu H, Welsh W, Yang CS. Tea polyphenol (-)-epigallocatechin-3-gallate inhibits DNA methyltransferase and reactivates methylation-silenced genes in cancer cell lines. *Cancer research* 2003; 63:7563-70.
54. Brueckner B, Garcia Boy R, Siedlecki P, Musch T, Kliem HC, Zielenkiewicz P, Suhai S, Wiessler M, Lyko F. Epigenetic reactivation of tumor suppressor genes by a novel small-molecule inhibitor of human DNA methyltransferases. *Cancer research* 2005; 65:6305-11.
55. Lind GE, Skotheim RI, Lothe RA. The epigenome of testicular germ cell tumors. *APMIS* 2007; 115:1147-60.
56. Netto GJ, Nakai Y, Nakayama M, Jadallah S, Toubaji A, Nonomura N, Albadine R, Hicks JL, Epstein JI, Yegnasubramanian S, et al. Global DNA hypomethylation in intratubular germ cell neoplasia and seminoma, but not in nonseminomatous male germ cell tumors. *Mod Pathol* 2008; 21:1337-44.
57. Wermann H, Stoop H, Gillis AJ, Honecker F, van Gurp RJ, Ammerpohl O, Richter J, Oosterhuis JW, Bokemeyer C, Looijenga LH. Global DNA methylation in fetal human germ cells and germ cell tumours: association with differentiation and cisplatin resistance. *J Pathol* 2010; 221:433-42.


58. Cerami E, Gao J, Dogrusoz U, Gross BE, Sumer SO, Aksoy BA, Jacobsen A, Byrne CJ, Heuer ML, Larsson E, et al. The cBio cancer genomics portal: an open platform for exploring multidimensional cancer genomics data. *Cancer discovery* 2012; 2:401-4.
59. Gao J, Aksoy BA, Dogrusoz U, Dresdner G, Gross B, Sumer SO, Sun Y, Jacobsen A, Sinha R, Larsson E, et al. Integrative analysis of complex cancer genomics and clinical profiles using the cBioPortal. *Sci Signal* 2013; 6:pl1.
60. Koul S, Houldsworth J, Mansukhani MM, Donadio A, McKiernan JM, Reuter VE, Bosl GJ, Chaganti RS, Murty VV. Characteristic promoter hypermethylation signatures in male germ cell tumors. *Mol Cancer* 2002; 1:8.
61. Koul S, McKiernan JM, Narayan G, Houldsworth J, Bacik J, Dobrzynski DL, Assaad AM, Mansukhani M, Reuter VE, Bosl GJ, et al. Role of promoter hypermethylation in Cisplatin treatment response of male germ cell tumors. *Mol Cancer* 2004; 3:16.
62. Beyrouthy MJ, Garner KM, Hever MP, Freemantle SJ, Eastman A, Dmitrovsky E, Spinella MJ. High DNA methyltransferase 3B expression mediates 5-aza-deoxycytidine hypersensitivity in testicular germ cell tumors. *Cancer research* 2009; 69:9360-6.
63. Candelaria M, Gallardo-Rincon D, Arce C, Cetina L, Aguilar-Ponce JL, Arrieta O, Gonzalez-Fierro A, Chavez-Blanco A, de la Cruz-Hernandez E, Camargo MF, et al. A phase II study of epigenetic therapy with hydralazine and magnesium valproate to overcome chemotherapy resistance in refractory solid tumors. *Annals of oncology : official journal of the European Society for Medical Oncology / ESMO* 2007; 18:1529-38.
64. Besaratinia A, Cockburn M, Tommasi S. Alterations of DNA methylome in human bladder cancer. *Epigenetics* 2013; 8:1013-22.
65. Veerla S, Panagopoulos I, Jin Y, Lindgren D, Hoglund M. Promoter analysis of epigenetically controlled genes in bladder cancer. *Genes Chromosomes Cancer* 2008; 47:368-78.
66. Maruyama R, Toyooka S, Toyooka KO, Harada K, Virmani AK, Zochbauer-Muller S, Farinas AJ, Vakar-Lopez F, Minna JD, Sagalowsky A, et al. Aberrant promoter methylation profile of bladder cancer and its relationship to clinicopathological features. *Cancer research* 2001; 61:8659-63.
67. Bilgrami SM, Qureshi SA, Pervez S, Abbas F. Promoter hypermethylation of tumor suppressor genes correlates with tumor grade and invasiveness in patients with urothelial bladder cancer. *Springerplus* 2014; 3:178.
68. Wang X, Chen E, Yang X, Wang Y, Quan Z, Wu X, Luo C. 5-azacytidine inhibits the proliferation of bladder cancer cells via reversal of the aberrant hypermethylation of the hepaCAM gene. *Oncol Rep* 2016; 35:1375-84.
69. Christoph F, Kempkensteffen C, Weikert S, Kollermann J, Krause H, Miller K, Schostak M, Schrader M. Methylation of tumour suppressor genes APAF-1 and DAPK-1 and in vitro effects of demethylating agents in bladder and kidney cancer. *British journal of cancer* 2006; 95:1701-7.
70. Hahn NM, Bonney PL, Dhawan D, Jones DR, Balch C, Guo Z, Hartman-Frey C, Fang F, Parker HG, Kwon EM, et al. Subcutaneous 5-azacitidine treatment of naturally occurring canine urothelial carcinoma: a novel epigenetic approach to human urothelial carcinoma drug development. *The Journal of urology* 2012; 187:302-9.
71. Xylinas E HM, Zhuang D, Krzywinski M, Erdem Z, Robinson BD, Elemento O, Clozel T, Shariat SF. An Epigenomic Approach to Improving Response to Neoadjuvant Cisplatin Chemotherapy in Bladder Cancer. *Biomolecules* 2016 Sep 2;6(3) pii: E37 doi: 103390/biom6030037.
72. Chuang JC, Warner SL, Vollmer D, Vankayalapati H, Redkar S, Bearss DJ, Qiu X, Yoo CB, Jones PA. S110, a 5-Aza-2'-deoxycytidine-containing dinucleotide, is an effective DNA methylation inhibitor in vivo and can reduce tumor growth. *Mol Cancer Ther* 2010; 9:1443-50.

73. Cheng JC, Weisenberger DJ, Gonzales FA, Liang G, Xu GL, Hu YG, Marquez VE, Jones PA. Continuous zebularine treatment effectively sustains demethylation in human bladder cancer cells. *Molecular and cellular biology* 2004; 24:1270-8.
74. Cheng JC, Matsen CB, Gonzales FA, Ye W, Greer S, Marquez VE, Jones PA, Selker EU. Inhibition of DNA methylation and reactivation of silenced genes by zebularine. *J Natl Cancer Inst* 2003; 95:399-409.
75. Ben-Kasus T, Ben-Zvi Z, Marquez VE, Kelley JA, Agbaria R. Metabolic activation of zebularine, a novel DNA methylation inhibitor, in human bladder carcinoma cells. *Biochem Pharmacol* 2005; 70:121-33.
76. Dote H, Cerna D, Burgan WE, Carter DJ, Cerra MA, Hollingshead MG, Camphausen K, Tofilon PJ. Enhancement of in vitro and in vivo tumor cell radiosensitivity by the DNA methylation inhibitor zebularine. *Clinical cancer research : an official journal of the American Association for Cancer Research* 2005; 11:4571-9.
77. Qi D, Li J, Que B, Su J, Li M, Zhang C, Yang M, Zhou G, Ji W. Long non-coding RNA DBCCR1-003 regulate the expression of DBCCR1 via DNMT1 in bladder cancer. *Cancer Cell Int* 2016; 16:81.
78. Habuchi T, Luscombe M, Elder PA, Knowles MA. Structure and methylation-based silencing of a gene (DBCCR1) within a candidate bladder cancer tumor suppressor region at 9q32-q33. *Genomics* 1998; 48:277-88.
79. Chuang JC, Warner SL, Vollmer D, Vankayalapati H, Redkar S, Bearss DJ, Qiu X, Yoo CB, Jones PA. S110, a 5-Aza-2'-Deoxycytidine-Containing Dinucleotide, Is an Effective DNA Methylation Inhibitor *In vivo* and Can Reduce Tumor Growth. *Molecular Cancer Therapeutics* 2010; 9:1443-50.
80. Lin J, Gilbert J, Rudek MA, Zwiebel JA, Gore S, Jiemjit A, Zhao M, Baker SD, Ambinder RF, Herman JG, et al. A phase I dose-finding study of 5-azacytidine in combination with sodium phenylbutyrate in patients with refractory solid tumors. *Clinical cancer research : an official journal of the American Association for Cancer Research* 2009; 15:6241-9.
81. Braithe F, Soriano AO, Garcia-Manero G, Hong D, Johnson MM, Silva Lde P, Yang H, Alexander S, Wolff J, Kurzrock R. Phase I study of epigenetic modulation with 5-azacytidine and valproic acid in patients with advanced cancers. *Clinical cancer research : an official journal of the American Association for Cancer Research* 2008; 14:6296-301.
82. Plummer R, Vidal L, Griffin M, Lesley M, de Bono J, Coulthard S, Sludden J, Siu LL, Chen EX, Oza AM, et al. Phase I study of MG98, an oligonucleotide antisense inhibitor of human DNA methyltransferase 1, given as a 7-day infusion in patients with advanced solid tumors. *Clinical cancer research : an official journal of the American Association for Cancer Research* 2009; 15:3177-83.
83. Kawakami T, Okamoto K, Ogawa O, Okada Y. Multipoint methylation and expression analysis of tumor suppressor genes in human renal cancer cells. *Urology* 2003; 61:226-30.
84. Avissar-Whiting M, Koestler DC, Houseman EA, Christensen BC, Kelsey KT, Marsit CJ. Polycomb group genes are targets of aberrant DNA methylation in renal cell carcinoma. *Epigenetics* 2011; 6:703-9.
85. Shenoy N, Vallumsetla N, Zou Y, Galeas JN, Shrivastava M, Hu C, Susztak K, Verma A. Role of DNA methylation in renal cell carcinoma. *J Hematol Oncol* 2015; 8:88.
86. Malouf GG, Su X, Zhang J, Creighton CJ, Ho TH, Lu Y, Raynal NJ, Karam JA, Tamboli P, Allanick F, et al. DNA Methylation Signature Reveals Cell Ontogeny of Renal Cell Carcinomas. *Clinical cancer research : an official journal of the American Association for Cancer Research* 2016; 22:6236-46.
87. Morris MR, Ricketts CJ, Gentle D, McDonald F, Carli N, Khalili H, Brown M, Kishida T, Yao M, Banks RE, et al. Genome-wide methylation analysis identifies epigenetically inactivated candidate tumour suppressor genes in renal cell carcinoma. *Oncogene* 2011; 30:1390-401.
88. Li M, Wang Y, Song Y, Bu R, Yin B, Fei X, Guo Q, Wu B. Aberrant DNA methyltransferase 1 expression in clear cell renal cell carcinoma development and progression. *Chin J Cancer Res* 2014; 26:371-81.

89. Peters I, Reese C, Dubrowskaja N, Antonopoulos WI, Krause M, Dang TN, Grote A, Becker A, Hennenlotter J, Stenzl A, et al. DNA methylation signature for the assessment of metastatic risk in primary renal cell cancer. *Journal of Clinical Oncology* 2017; 35:516-516.
90. Ricketts CJ, Morris MR, Gentle D, Shuib S, Brown M, Clarke N, Wei W, Nathan P, Latif F, Maher ER. Methylation profiling and evaluation of demethylating therapy in renal cell carcinoma. *Clin Epigenetics* 2013; 5:16.
91. Alkamal I, Ikromov O, Tolle A, Fuller TF, Magheli A, Miller K, Krause H, Kempkensteffen C. An epigenetic screen unmasks metallothioneins as putative contributors to renal cell carcinogenesis. *Urol Int* 2015; 94:99-110.
92. Alleman WG, Tabios RL, Chandramouli GV, Aprelikova ON, Torres-Cabala C, Mendoza A, Rogers C, Sopko NA, Linehan WM, Vasselli JR. The in vitro and in vivo effects of re-expressing methylated von Hippel-Lindau tumor suppressor gene in clear cell renal carcinoma with 5-aza-2'-deoxycytidine. *Clinical cancer research : an official journal of the American Association for Cancer Research* 2004; 10:7011-21.
93. Coral S, Parisi G, Nicolay HJ, Colizzi F, Danielli R, Fratta E, Covre A, Taverna P, Sigalotti L, Maio M. Immunomodulatory activity of SGI-110, a 5-aza-2'-deoxycytidine-containing demethylating dinucleotide. *Cancer immunology, immunotherapy : CII* 2013; 62:605-14.
94. Reu FJ, Bae SI, Cherkassky L, Leaman DW, Lindner D, Beaulieu N, MacLeod AR, Borden EC. Overcoming resistance to interferon-induced apoptosis of renal carcinoma and melanoma cells by DNA demethylation. *Journal of clinical oncology : official journal of the American Society of Clinical Oncology* 2006; 24:3771-9.
95. Chiam K, Centenera MM, Butler LM, Tilley WD, Bianco-Miotto T. GSTP1 DNA methylation and expression status is indicative of 5-aza-2'-deoxycytidine efficacy in human prostate cancer cells. *PloS one* 2011; 6:e25634.
96. Ikromov O, Alkamal I, Magheli A, Ratert N, Sendeski M, Miller K, Krause H, Kempkensteffen C. Functional Epigenetic Analysis of Prostate Carcinoma: A Role for Seryl-tRNA Synthetase? *J Biomark* 2014; 2014:362164.
97. Sabatino MA, Geroni C, Ganzinelli M, Ceruti R, Broggin M. Zebularine partially reverses GST methylation in prostate cancer cells and restores sensitivity to the DNA minor groove binder brostallicin. *Epigenetics* 2013; 8:656-65.
98. Collard RL, Harya NS, Monzon FA, Maier CE, O'Keefe DS. Methylation of the ASC gene promoter is associated with aggressive prostate cancer. *The Prostate* 2006; 66:687-95.
99. Gertych A, Oh JH, Wawrowsky KA, Weisenberger DJ, Tajbakhsh J. 3-D DNA methylation phenotypes correlate with cytotoxicity levels in prostate and liver cancer cell models. *BMC Pharmacol Toxicol* 2013; 14:11.
100. Kim TD, Jin F, Shin S, Oh S, Lightfoot SA, Grande JP, Johnson AJ, van Deursen JM, Wren JD, Janknecht R. Histone demethylase JMJD2A drives prostate tumorigenesis through transcription factor ETV1. *J Clin Invest* 2016; 126:706-20.
101. Festuccia C, Gravina GL, D'Alessandro AM, Muzi P, Millimaggi D, Dolo V, Ricevuto E, Vicentini C, Bologna M. Azacitidine improves antitumor effects of docetaxel and cisplatin in aggressive prostate cancer models. *Endocr Relat Cancer* 2009; 16:401-13.
102. Frost P, Abbruzzese JL, Hunt B, Lee D, Ellis M. Synergistic cytotoxicity using 2'-deoxy-5-azacytidine and cisplatin or 4-hydroperoxycyclophosphamide with human tumor cells. *Cancer Res* 1990; 50:4572-7.
103. Kagey JD, Kapoor-Vazirani P, McCabe MT, Powell DR, Vertino PM. Long-term stability of demethylation after transient exposure to 5-aza-2'-deoxycytidine correlates with sustained RNA polymerase II occupancy. *Mol Cancer Res* 2010; 8:1048-59.
104. McCabe MT, Low JA, Daignault S, Imperiale MJ, Wojno KJ, Day ML. Inhibition of DNA methyltransferase activity prevents tumorigenesis in a mouse model of prostate cancer. *Cancer research* 2006; 66:385-92.

105. Sonpavde G, Aparicio AM, Zhan F, North B, Delaune R, Garbo LE, Rousey SR, Weinstein RE, Xiao L, Boehm KA, et al. Azacitidine favorably modulates PSA kinetics correlating with plasma DNA LINE-1 hypomethylation in men with chemonaive castration-resistant prostate cancer. *Urologic oncology* 2011; 29:682-9.
106. Singal R, Ramachandran K, Gordian E, Quintero C, Zhao W, Reis IM. Phase I/II study of azacitidine, docetaxel, and prednisone in patients with metastatic castration-resistant prostate cancer previously treated with docetaxel-based therapy. *Clinical genitourinary cancer* 2015; 13:22-31.
107. Rudek MA, Zhao M, He P, Hartke C, Gilbert J, Gore SD, Carducci MA, Baker SD. Pharmacokinetics of 5-azacitidine administered with phenylbutyrate in patients with refractory solid tumors or hematologic malignancies. *Journal of clinical oncology : official journal of the American Society of Clinical Oncology* 2005; 23:3906-11.
108. Wong CM, Wong CC, Ng YL, Au SL, Ko FC, Ng IO. Transcriptional repressive H3K9 and H3K27 methylations contribute to DNMT1-mediated DNA methylation recovery. *PLoS One* 2011; 6:e16702.
109. Kuasne H, Marchi FA, Rogatto SR, de Syllos Colus IM. Epigenetic mechanisms in penile carcinoma. *Int J Mol Sci* 2013; 14:10791-808.
110. Feber A, Arya M, de Winter P, Saqib M, Nigam R, Malone PR, Tan WS, Rodney S, Lechner M, Freeman A, et al. Epigenetics markers of metastasis and HPV-induced tumorigenesis in penile cancer. *Clinical cancer research : an official journal of the American Association for Cancer Research* 2015; 21:1196-206.
111. Kuasne H, Colus IM, Busso AF, Hernandez-Vargas H, Barros-Filho MC, Marchi FA, Scapulatempo-Neto C, Faria EF, Lopes A, Guimaraes GC, et al. Genome-wide methylation and transcriptome analysis in penile carcinoma: uncovering new molecular markers. *Clin Epigenetics* 2015; 7:46.

Figure 1. Schematic overview of the literature search.



Key words: epigenetics, DNMT, DNMTi, genitourinary, testicular cancer, germ-cell tumors, bladder cancer, renal cell carcinoma, prostate cancer, penile cancer, azacitidine, decitabine, guadecitabine, zebularine, non-nucleoside


Abbreviations: ASCO – American Society of Clinical Oncology, ASCO GU – Genitourinary Cancers Symposium, AACR – American Association for Cancer Research, GCTs – germ cell tumors, RCC – renal cell carcinoma


Table 1. Inclusion, exclusion criteria and the selection process for including publications inclusion in the review article

<i>Inclusion criteria</i>	<i>Exclusion criteria</i>
<ol style="list-style-type: none"> 1. Pre-clinical <i>in vitro</i> studies evaluating DNA methyltransferase inhibitors (DNMTis) in genitourinary (GU) tumors including germ-cell tumors, bladder cancer, renal cell carcinoma, prostate cancer and penile cancer. 2. Pre-clinical <i>in vivo</i> studies evaluating DNMTis in GU tumors including germ-cell tumors, bladder cancer, renal cell carcinoma, prostate cancer and penile cancer. 	<ol style="list-style-type: none"> 1. Title including diagnoses, cell lines and animal models for cancers different than GU (e.g. hematologic malignancies, other solid tumors). 2. Abstract including cell lines, animal models and clinical trials for cancers different than GU. 3. Full text articles including cell lines, animal models and clinical trials for cancers other than GU tumors.

3. Clinical studies evaluating DNMTis in GU tumors including germ-cell tumors, bladder cancer, renal cell carcinoma, prostate cancer and penile cancer.	
<p>Selection process of published works was conducted in several steps. First, review of publication titles retrieved from the literature search was performed. If title met the inclusion criteria, the abstract was reviewed. If abstract met the inclusion criteria, the full text article was reviewed. If inclusion criteria for full text article were met, the article was included in our review manuscript. If title of general nature was found (e.g. DNMTis in cancer, solid tumors etc.), the abstract was reviewed. Subsequently, if abstract did not clearly specify the cancer types, the full text article was reviewed and included or excluded according to the inclusion/exclusion criteria.</p>	

Figure 2. The expression of DNMT3A/B in different types of tumors ^{58, 59}





Abbreviations: DLBCL – diffuse large B-cell lymphoma, chRCC – chromophobe renal cell cancer, PCPG – pheochromocytoma and paraganglioma, ACC – adrenocortical carcinomas, ccRCC – clear cell renal cell carcinoma, pRCC – papillary renal cell carcinoma, AML – acute myeloid leukemia, GBM - glioblastoma

Table 2. Ongoing clinical studies with DNMTis listed on *clinicaltrials.gov*

<i>Study ID</i>	<i>Phase</i>	<i>Treatments</i>	<i>Indication</i>	<i>Planned patients (N)</i>	<i>Date started</i>	<i>Planned completion date</i>
NCT02429466	I	SGL-110 Plus Cisplatin	Relapsed Platinum Refractory Germ Cell Tumors	15	April,2015	May 2018
NCT02223052	I	Oral Azacitidine	Adult Cancer Subjects including genitourinary cancers	60	October,2017	January 2018
NCT02788201	I/II	75 Approved Agents	Advanced Urothelial Carcinoma	20	May 20, 2016	June 29, 2019
NCT02961101	I/II	Anti-PD-1 antibody plus Decitabine	Relapsed or Refractory Malignancies Including RCC	100	May 2016	May 2020

NCT01799083	I/II	Decitabine Alone and/or in Combination with Chemotherapy and/or Cytokine Induced Killer Cell Transfusion	Relapsed or Refractory Solid Tumors and B Cell Lymphomas	100	December 2012	December 2017
NCT02423057	I	4'-Thio-2'-Deoxycytidine	Advanced Solid Tumors	46	April 13, 2015	April 18, 2018
NCT00978250	II	5-Fluoro-2'-Deoxycytidine with Tetrahydrouridine	Solid tumors including urinary bladder neoplasms	165	August 20, 2009	May 27, 2017
NCT02959437	I/II	Azacitidine plus Pembrolizumab and Epacadostat	Advanced Solid Tumors	142	January 24, 2017	October 2021
NCT02998567	I	Guadecitabine plus Pembrolizumab	Castration-Resistant Prostatic Cancer and Non-Small Cell Lung Cancer	35	December 2016	November 2018
NCT02030067	I/II	Single Agent RX-3117	Metastatic Bladder and Pancreatic Cancer	72	December 2013	December 2017